

Non-strictly-proper transforms section 2.2.5, p.16 in Bodson text

What if the order of the numerator is equal to or even greater than the order of the denominator? $m \geq n$?

Example: $F(s) = \frac{2 \cdot s^2 + 100}{s^2 + 8 \cdot s + 41}$ $m := 2$
 $n := 2$

First divide, before partial fraction expansion $s^2 + 8 \cdot s + 41 \left\{ \begin{array}{l} \overline{2 \cdot s^2 + 0 \cdot s + 100} \\ \hline \end{array} \right.$

"remainder"

$$F(s) = \frac{2 \cdot s^2 + 100}{s^2 + 8 \cdot s + 41} =$$

$$f(t) =$$

Delta functions are not very common in real life.

Non-strictly-proper transforms are just as common.

Properties of Signals

Can you tell what $f(t)$ must be just by looking at $F(s)$? YES, somewhat...

$$\frac{s+5}{s \cdot (s^2 + 4 \cdot s + 13) \cdot (s-10)}$$

$$\frac{s+5}{s \cdot (s^2 + 64) \cdot (s+10)}$$

$$\frac{s+5}{s \cdot (s^2 - 4 \cdot s + 13) \cdot (s+10)}$$

$$\frac{s+5}{s \cdot (s^2 + 4 \cdot s + 13)^2 \cdot (s+10)}$$

$$\frac{s+5}{s^3 \cdot (s^2 + 4 \cdot s + 13)^2 \cdot (s+10)^2}$$